

A NOTE ON THE UNIFORMIZATION OF GRADIENT KÄHLER RICCI SOLITONS

ALBERT CHAU¹ AND LUEN-FAI TAM²

ABSTRACT. Applying a well known result for attracting fixed points of biholomorphisms [4, 6], we observe that one immediately obtains the following result: if (M^n, g) is a complete non-compact gradient Kähler-Ricci soliton which is either steady with positive Ricci curvature so that the scalar curvature attains its maximum at some point, or expanding with non-negative Ricci curvature, then M is biholomorphic to $\mathbb{C}^n$.

We will show the following:

Theorem 1. *If (M^n, g) is a complete non-compact gradient Kähler-Ricci soliton which is either steady with positive Ricci curvature so that the scalar curvature attains its maximum at some point, or expanding with non-negative Ricci curvature, then M is biholomorphic to $\mathbb{C}^n$.*

Recall that a Kähler manifold $(M, g_{i\bar{j}}(x))$ is said to be a Kähler-Ricci soliton if there is a family of biholomorphisms ϕ_t on M , given by a holomorphic vector field V , such that $g_{ij}(x, t) = \phi_t^*(g_{ij}(x))$ is a solution of the Kähler-Ricci flow:

$$(0.1) \quad \begin{aligned} \frac{\partial}{\partial t} g_{i\bar{j}} &= -R_{i\bar{j}} - 2\rho g_{i\bar{j}} \\ g_{i\bar{j}}(x, 0) &= g_{i\bar{j}}(x) \end{aligned}$$

for $0 \leq t < \infty$, where $R_{i\bar{j}}$ denotes the Ricci tensor at time t and ρ is a constant. If $\rho = 0$, then the Kähler-Ricci soliton is said to be of *steady type* and if $\rho > 0$ then the Kähler-Ricci soliton is said to be of *expanding type*. We always assume that g is complete and M is non-compact. If in addition, the holomorphic vector field is given by the gradient of a real valued function f , then it is called a gradient Kähler-Ricci soliton.

Date: April 2004 (Revised in July 2004).

2000 *Mathematics Subject Classification.* Primary 53C44; Secondary 58J37, 35B35.

¹Research partially supported by The Institute of Mathematical Sciences, The Chinese University of Hong Kong, Shatin, Hong Kong, China.

²Research partially supported by Earmarked Grant of Hong Kong #CUHK4032/02P.

Note that in this case, we have that

$$(0.2) \quad \begin{aligned} f_{i\bar{j}} &= R_{i\bar{j}} + 2\rho g_{i\bar{j}} \\ f_{ij} &= 0. \end{aligned}$$

If (M, g) is a gradient Kähler-Ricci soliton (of steady or expanding type) which is either steady with positive Ricci curvature so that the scalar curvature attains its maximum at some point, or expanding with non-negative Ricci curvature, then one can show that ϕ_t , the flow on M along the vector field ∇f , satisfies:

- (i) ϕ_t is a biholomorphism from M to M for all $t \geq 0$,
- (ii) ϕ_t has a unique fixed point p , i.e. $\phi_t(p) = p$ for all $t \geq 0$,
- (iii) M is attracted to p under ϕ_t in the sense that for any open neighborhood U of p and for any compact subset W of M , there exists $T > 0$ such that $\phi_t(W) \subset U$ for all $t \geq T$.

Condition (i) is clear. Condition (ii) is shown in [2, 3]. To see that condition (iii) holds, we consider any $R > 0$ and let $B(R)$ be the geodesic ball of radius R with center at p with respect to the metric $g(0)$. From the proof of Lemma 3.2 in [2], there exists $C_R > 0$ such that for any $q \in B(R)$ and for any $v \in T^{1,0}(M)$ at q ,

$$\|v\|_{\phi_t^*(g)} \leq \exp(-C_R t) \|v\|_g.$$

Since $\phi_t(p) = p$, it is easy to see that given any open set $U \subset M$ containing p , we have $\phi_t(B(R)) \subset U$ provided t is large, and thus condition (iii) is satisfied.

The following theorem was proved for the case $M = \mathbb{C}^n$ in [4], and was later observed to be true on a general complex manifold M in [6].

Theorem 2. *Let F be a biholomorphism from a complex manifold M^n to itself and let $p \in M^n$ be a fixed point for F . Fix a complete Riemannian metric g on M and define*

$$\Omega := \{x \in M : \lim_{k \rightarrow \infty} \text{dist}_g(F^k(x), p) = 0\}$$

where $F^k = F \circ F^{k-1}$, $F^1 = F$. Then Ω is biholomorphic to $\mathbb{C}^n$ provided Ω contains an open neighborhood around p .

Proof of Theorem 1. By conditions (i)-(iii) we may apply Theorem 2 to the biholomorphism $\phi_1 : M \rightarrow M$ to conclude that M is biholomorphic to $\mathbb{C}^n$. $\square$

Remark 1. In the first version of this article we proved Theorem 2 in a special case. We would like to thank Dror Varolin for pointing out to us that what we proved had been known earlier [4, 6].

Remark 2. After posting the first version of this article we learned that Theorem 1 in the case of a steady gradient Kähler Ricci soliton had been known independently to Robert Bryant [1].

REFERENCES

1. Bryant, Robert, *Gradient Kahler Ricci Solitons*, arXiv eprint 2004. arXiv:math.DG/0407453
2. Chau, Albert and Tam, Luen-fai, *Gradient Kähler-Ricci soliton and a uniformization conjecture*, arXiv eprint 2003. arXiv:math.DG/0310198.
3. Hamilton, Richard S., *Formation of Singularities in the Ricci Flow*, Contemporary Mathematics, **71** (1988), 237-261.
4. Rosay, J.P. and Rudin, W., *Holomorphic Maps from $\mathbb{C}^n$ to $\mathbb{C}^n$* , Trans. AMS **310** (1988), 47-86.
5. Sternberg, Shlomo, *Local contractions and a theorem of Poincaré*, Amer. J. Math., **79** (1957), 809–824.
6. Varolin, Dror, *The density property for complex manifolds and geometric structures II*, Internat. J. Math. **11** (2000), no. 6, 837–847.

HARVARD UNIVERSITY, DEPARTMENT OF MATHEMATICS, ONE OXFORD STREET,
CAMBRIDGE, MA 02138, USA

E-mail address: `chau@math.harvard.edu`

DEPARTMENT OF MATHEMATICS, THE CHINESE UNIVERSITY OF HONG KONG,
SHATIN, HONG KONG, CHINA.

E-mail address: `lftam@math.cuhk.edu.hk`